


New resonances in B-meson decays

R. CHISTOV

(for the Belle Collaboration)

Institute for Theoretical and Experimental Physics, Moscow, Russia



The $X(3872)$ and $Y(3940)$ properties and decay modes from Belle are reviewed and the results on a search for the decay $B^+ \rightarrow h_c K^+$, $h_c \rightarrow \eta_c \gamma$ at Belle are presented.

1 Introduction

B -mesons have proved to be a rich source of new particles. The B-factories at KEK (Belle) and SLAC (BaBar) extensively use the exclusive production of J/ψ , χ_{c1} and η_c charmonia for the CP violation measurements. In addition to these conventional charmonia Belle observed the production of $\eta_c(2S)$ ¹ and $\psi(3770)$ ² in exclusive $B \rightarrow (c\bar{c})K^+$ decays. Belle and BaBar also observed inclusive χ_{c2} production in B decays ³. In 2003, by analyzing the $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$ decays, Belle observed a narrow charmonium-like new state (denoted as $X(3872)$) decaying into $J/\psi \pi^+ \pi^-$ ⁴. Recently Belle reported the observation of $Y(3940) \rightarrow J/\psi \omega$ in $B^+ \rightarrow Y(3940) K^+$ decays ⁵.

All observed resonances – from $\eta_c(2S)$ to $Y(3940)$ – are difficult to reconstruct without the constraints provided by B decays, as they decay to high-multiplicity final states. B mesons also provide an excellent opportunity to test different hypotheses for the J^P quantum numbers of these resonances via decay angle analysis.

2 Observation of $X(3872)$ and its properties

Just after the discovery of $X(3872)$ by Belle, this new state was confirmed by the CDF, D0 and BaBar collaborations ⁶. Its mass was measured to be 3871.9 ± 0.5 MeV which is very close to the $D^0 \bar{D}^{*0}$ threshold of 3871.3 ± 1.0 MeV. All these first measurements directly provide a lot of information on the $X(3872)$ properties ⁴:




Figure 1: The invariant mass spectrum of $\pi^+ \pi^-$ pairs from the $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ decay.

- the two pion mass from the $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ decay tends to peak near the ρ^0 mass, consistent with positive C-parity of the $X(3872)$;
- the decays $X(3872) \rightarrow \chi_{c1,2} \gamma$ are not seen; this likely excludes the $1^3D_{2,3}$ ($\psi_{2,3}$) assignment for the $X(3872)$;
- the decay $X(3872) \rightarrow D\bar{D}$ is suppressed or forbidden²; this, together with its narrow width ($\Gamma < 2.7$ MeV), suggests $J^P = 0^+, 1^-, 2^+, \dots$ are ruled out;

CDF and D0 have measured the $X(3872)$ production properties to be very similar to those of the $\psi(2S)$ ⁶. BaBar reported a null search for the charged $X(3872)$ partners in $B \rightarrow J/\psi \pi^+ \pi^0 K^+$ decays⁷ and a null search for $X(3872) \rightarrow J/\psi \eta$ decay⁸. The former rules out the isovector hypothesis and the latter excludes the presence of gluonic degrees of freedom in the $X(3872)$ wave function.

Fig. 1 shows the $M(\pi^+ \pi^-)$ spectrum from the updated analysis of $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ decays by Belle (253 fb^{-1}). The ρ^0 signal is strong and supports $C(X(3872)) = +1$.

Recently Belle has found evidence for another decay mode of $X(3872)$: $X(3872) \rightarrow J/\psi \omega^*$ where ω^* is virtual and reconstructed in the $\pi^+ \pi^- \pi^0$ channel. According to Swanson's model⁹, this observation supports the $D^0 \bar{D}^{*0}$ molecular interpretation of $X(3872)$. The full set of the most recent Belle results on the $X(3872)$ properties can be found elsewhere¹⁰.

3 Observation of $Y(3940)$

By analyzing exclusive $B^+ \rightarrow J/\psi \pi^+ \pi^- \pi^0 K^+$ decays Belle observed a new resonance $Y(3940)$ decaying to $J/\psi \omega$ ⁵. Fig. 2 shows the $M(J/\psi \omega)$ distribution for the B -meson candidates. The curve in Fig. 2 (a) indicates the result of a fit with only a phase-space-like two-body threshold function. The curve in Fig. 2 (b) shows the result of a fit that includes an S-wave Breit-Wigner resonance term. The mass and width of $Y(3940)$ were measured to be $3943 \pm 11 \pm 13$ MeV and $87 \pm 22 \pm 26$ MeV, respectively. The observed state is above the $D\bar{D}^{(*)}$ threshold and would decay predominantly to $D\bar{D}$ and/or $D\bar{D}^*$ if it is a $c\bar{c}$ charmonium. In contrast, for a $c\bar{c}$ - gluon hybrid the open charm decay modes are suppressed or forbidden. So the observed $Y(3940)$ is a possible candidate for the first $c\bar{c}$ - gluon hybrid state.

4 Search for $B^+ \rightarrow h_c K^+$

The 1^1P_1 , $J^P = 1^{+-}$ h_c has for a long time been a missing state. Recently CLEO reported the observation of h_c in $\psi(2S) \rightarrow h_c \pi^0$, $h_c \rightarrow \eta_c \gamma$ decays¹¹. The mass was measured to be $3524.4 \pm 0.6 \pm 0.4$ MeV - in agreement with theoretical expectations that the $M(h_c)$ is close to the c.o.g. of the $< 1^3P_J >$ triplet.




Figure 2: The invariant mass spectrum of $J/\psi \omega$ combinations from the $B^+ \rightarrow J/\psi \omega K^+$ decay. The curve in (a) indicates the result of a fit with only a phase-space-like two-body threshold function. The curve in (b) shows the result of a fit that includes an S-wave Breit-Wigner resonance term.

Belle searched for h_c production in exclusive $B^+ \rightarrow h_c K^+$, $h_c \rightarrow \eta_c \gamma$ decays. Fig. 3 shows the $M(\eta_c \gamma)$ for the B -candidates. No evidence for a signal around the CLEO h_c mass is seen. Belle set an upper limit for $\mathcal{B}(B^+ \rightarrow h_c K^+) \times \mathcal{B}(h_c \rightarrow \eta_c \gamma)$ that is less than 1.5×10^{-4} for $M(h_c) \sim 3520$ MeV¹².

5 Study of $D_{sJ}(2317)$ and $D_{sJ}(2460)$

The $D_{sJ}(2317)^{*+}$ and $D_{sJ}(2460)^+$ mesons were observed by BaBar¹³ and CLEO¹⁴. Belle confirmed these states and observed their production in exclusive $B \rightarrow D_{sJ}^{*+} \bar{D}^{(*)}$ decays¹⁵. The observation of these decays allowed us to perform decay angle analysis. Belle data support $J = 0$ for $D_{sJ}(2317)$ and $J = 1$ for $D_{sJ}(2460)$ ¹⁶.

6 Summary

B mesons provide a clean environment for the observation of yet-unseen charmonia and other new unexpected resonances and the understanding of their properties.

The nature of new resonances $X(3872)$ and $Y(3940)$ remains unclear so far. They could be either $c\bar{c}$ states or exotic hadrons: $D^0 \bar{D}^{*0}$ molecular ($X(3872)$) and $c\bar{c} - gluon$ hybrid ($Y(3940)$).

Acknowledgments

We thank the KEKB group for the excellent operation of the accelerator, the KEK cryogenics group for the efficient operation of the solenoid, and the KEK computer group and the NII for valuable computing and Super-SINET network support. We acknowledge support from MEXT and JSPS (Japan); ARC and DEST (Australia); NSFC (contract No. 10175071, China); DST (India); the BK21 program of MOEHRD and the CHEP SRC program of KOSEF (Korea); KBN (contract No. 2P03B 01324, Poland); MIST (Russia); MHEST (Slovenia); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE (USA).




Figure 3: The invariant mass spectrum of $\eta_c \gamma$ for the $B^+ \rightarrow \eta_c \gamma K^+$ candidates.

References

1. Belle Collaboration, S.-K.Choi, S.L.Olsen, et al. *et al.*, Phys. Rev. Lett. **89**, 102001 (2002)
2. Belle Collaboration, R.Chistov, *et al.*, Phys. Rev. Lett. **93**, 051803 (2004)
3. Belle Collaboration, K.Abe, *et al.*, Phys. Rev. Lett. **89**, 011803 (2002); BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. D **67**, 032002 (2003)
4. Belle Collaboration, S.-K.Choi, S.L.Olsen, et al. *et al.*, Phys. Rev. Lett. **91**, 262001 (2003)
5. Belle Collaboration, S.-K.Choi, S.L.Olsen, et al. *et al.*, Phys. Rev. Lett. **94**, 182002 (2005)
6. CDF-II Collaboration, D. Acosta *et al.*, Phys. Rev. Lett. **93**, 072001 (2004); D0 Collaboration, V.M. Abazov *et al.*, Phys. Rev. Lett. **93**, 162002 (2004); BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. D **71**, 071103 (2005)
7. BaBar Collaboration, B. Aubert *et al.*, hep-ex/0412051
8. BaBar Collaboration, B. Aubert *et al.*, hep-ex/0402025
9. E. S. Swanson, Phys. Lett. B **589**, 189 (2004)
10. Belle Collaboration, K.Abe, *et al.*, hep-ex/0505038
11. CLEO Collaboration, J. L. Rosner, *et al.*, hep-ex/0505073, submitted to Phys. Rev. Lett.
12. Belle Collaboration, K.Abe, *et al.*, BELLE-CONF-0440
13. BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **90**, 242001 (2003)
14. CLEO Collaboration, D.Besson *et al.*, Phys. Rev. D **68** 032002, (2003)
15. Belle Collaboration, P.Krokovny, A.Bondar, *et al.*, Phys. Rev. Lett. **91**, 262002 (2003)
16. Belle Collaboration, K.Abe, *et al.*, BELLE-CONF-0461